

STATUS REPORT - PINNIPED PREDATION AND DETERRENT ACTIVITIES AT BONNEVILLE DAM, 2013

Robert Stansell, Bjorn van der Leeuw, and Karrie Gibbons - (541) 374-8801

Fisheries Field Unit
U.S. Army Corps of Engineers
Bonneville Lock and Dam
Cascade Locks, OR 97014

February 14, 2013

This is the second weekly status report for 2013 and summarizes all pinniped predation monitoring and deterrent activities at Bonneville Dam from January 1 through February 13, 2013 (unless otherwise noted). Regular daylight observations began January 4 and will continue through the end of May, five days per week excluding holidays. Final predation estimates will be expanded for hours and days not observed, adjusted for “unknown” prey species take, and a night time predation factor applies at the end of the observation season and those updated figures will be presented in our annual field report. This report can be found at: <http://www.nwd-wc.usace.army.mil/tmt/documents/fish/>

PRELIMINARY RESULTS

All data presented here are preliminary as of the status report date. Predation figures are unexpanded (unless otherwise noted) and sea lion abundance estimates will likely change as the season progresses and data are proofed and analyzed, so please use these estimates with appropriate caution. A final report summarizing the results of the 2013 evaluation will be available in the fall of this year.

PINNIPED ABUNDANCE

California sea lions (*Zalophus californianus* – CSL) have remained scarce so far this year (Figure 1), with only one observed on three separate days. We were unable to identify the CSL, and no predation was observed by any CSL yet. Steller sea lions (*Eumetopias jubatus* - SSL) are slightly less abundant than the past two years (figures 1 and 2). The maximum number of SSL seen any day so far this year was 21 and only 1 for CSL (Figure 1). We have documented approximately 30 different SSL's visiting the dam so far. All of the SSL are confirmed as seen in past years, although there may be some new individuals we have not confirmed yet.

A few SSL have been observed hauling out inside the end of the corner collector outfall, but mostly they are resting in the water in large pods in the spillway. The traps have been closed since early January.

PREDATION DATA

Unexpanded numbers for fish observed taken in the Bonneville Dam tailrace for 2013 (through February 13) are:

<i>Prey</i>	<i>California Sea Lions</i>	<i>Steller Sea Lions</i>	<i>Total</i>
Chinook	0	1	1
Steelhead	0	23	23
Sturgeon	0	244	244
Lamprey	0	4	4
Shad	0	65	65
Smolt	0	0	0
Other	0	0	0
Unknown	0	48	48

It is likely that most of the unknown fish caught by Steller sea lions were sturgeon. The Steller sea lions are catching many of the fish at the downstream range of our viewing area, making fish identification very difficult. Most sturgeon have been caught in the spillway tailrace, followed by the powerhouse 2 tailrace, while relatively few are caught at powerhouse 1 (Figure 3). Sturgeon catch (398 expanded for weekends only) is less than the previous four years at this time (Figure 4), most taken being in the 2 to 4 foot range. Few fish are passing the count stations (255 steelhead and 0 Chinook) from January 1 through February 12, lower than the last three years. Total salmonid catch through February 13 (37 expanded by interpolating for weekends) is fewer than the past six years (2012 – 93, 2011 – 85, 2010 – 103, 2009 – 65, 2008 – 66, 2007 - 96), all by Steller sea lions.

DETERRENTS/TRAPPING

Hazing by USDA (land) will begin on March 1 and will continue until the end of May, seven days a week. CRITFC will begin hazing from boats March 4.

The states have not conducted any trapping and do not plan to until early March due to funding issues.

OTHER ITEMS OF INTEREST

Nothing to report here from the past week.

Figure 1. Daily pinniped abundance, by species, at Bonneville Dam, 2013.




Figure 2. Average daily presence of pinnipeds, by species, to date (February 13) for each year at Bonneville Dam.




Figure 3. Distribution of prey taken by Pinnipeds by location at Bonneville Dam, through February 13, 2013.




Figure 4. Daily cumulative sturgeon catch (interpolated for weekends) at Bonneville Dam, 2006-2013.

