

U.S. Fish and Wildlife Service
SYSTEM OPERATIONAL REQUEST: #2008-FWS-3

TO: BG William E. Rapp	COE-NWD
James R. Adams	COE-NWD
Kevin Grode	COE-RCC
James D. Barton	COE-Water Management
Carolyn Fitzgerald	COE-NWS – Water Management
David Ponganis	COE-NWD
Col. Steven Wright	COE- NWS
Stephen J. Wright	BPA-Administrator
Steve Oliver	BPA-PG-5

FROM: Rich Torquemada, Acting Supervisor, Upper Columbia Fish and Wildlife Office, U.S. Fish and Wildlife Service, on behalf of the Kootenai Valley Resource Initiative.

DATE: 29 October 2008

SUBJECT: Winter Temperature Operation of Libby Dam for Kootenai River burbot

SPECIFICATIONS:

U.S. Fish and Wildlife Service staff, in cooperation with the Kootenai Valley Resource Initiative (KVRI) and as recommended by the KVRI burbot sub-committee, request that the U.S. Army Corps of Engineers use the selective withdrawal gate system at Libby Dam to release the coolest water possible in November and December, 2008, before temperature stratification limits the temperature control capability. The purpose of this operation is to provide cooler river temperatures downstream of Bonners Ferry, and also to determine how radio-tagged burbot in the Kootenai River respond to these temperatures. This will likely result in November and December temperatures slightly cooler than the existing selective withdrawal temperature rule curve (Figure 1). This deviation from the temperature rule curve has been coordinated with Montana Fish, Wildlife & Parks (MFWP). The Corps at Libby Dam will remove selective withdrawal gates incrementally during late October to assure that daily temperature change remains within 2 degrees F per day; gates will be removed systematically to slowly lower river temperature to the minimum by early November (on average, a span of about 5 degrees C, or 9 degrees F; Figure 1).




Figure 1. Selective withdrawal temperature guidelines for the Kootenai River below Libby Dam. The mean average release temperature was within the guidelines for the period displayed, with means slightly less than optimal early in the winter, and slightly higher than optimal later in the winter. There is water much cooler than optimal available for release during the early winter period until the reservoir becomes isothermic about mid-December (Figure 2).




Figure 2 – 2007 Libby Dam forebay temperature profile. The reservoir becomes isothermic by mid-December during most years.

JUSTIFICATION:

The request is designed to cool the river (Bonners Ferry vicinity) during the burbot migration period (November to mid-December) and to the extent possible during spawning season (Dec 15 – end of Feb), when temperatures of 1 to 4° C are preferable. Burbot spawning migrations may be affected by water temperature conditions much earlier in the winter/late autumn (Paragamian 2005, pers. comm.), particularly when combined with higher flows.

Lengthy migrations have been documented in the late fall/early winter and again in late winter/early spring that coincide with spawning (Robins and Deubler 1955, McCrimmon 1959, Percy 1975, Morrow 1980, Johnson 1981, Breeser et al. 1988, Evenson 2000, Paragamian 2000, Schram 2000). These migrations were often temporally correlated with changes in water temperatures, although movement appeared to be minimal immediately prior to spawning (Evenson 2000).

The following excerpts of actions to be implemented are from the widely agreed upon Kootenai River burbot conservation strategy (KVRI Burbot Committee, 2005):

9.4 Hydro Operations

9.4.1 Develop an experimental Kootenai River flow/water temperature operation to evaluate the effectiveness of restoring natural spawning and recruitment by reducing winter temperatures and velocities. Implement experimental operations when conditions allow to evaluate burbot spawning requirements while preserving flexibility in needed hydropower production and flood control operations. Annual operations will be coordinated through the Regional Forum Technical Management Team (TMT). The KVRI Burbot Committee will coordinate with the U.S. Fish and Wildlife Service to develop System Operations Requests (SOR) to the TMT to request flow conditions or temperature requirements in any given year.

9.4.4 Evaluate use of selective withdrawal during migratory pre-spawning periods to affect thermograph at Bonners and downstream to benefit burbot, and monitor water temperature at Porthill.

The principal migration monitoring method involved implanting radio transmitters in larger burbot, and then tracking their behavior with an array of fixed, continuously recording receivers along the Kootenai (y) River and at some tributary mouths. Seven burbot were implanted with *Vemco* coded transmitter tags between October 2005 and March 2006. Three were tagged near Nick's Island (rkm 144.5), two were tagged near the mouth of the Goat River (rkm 152.7) and two were tagged at Ambush Rock (rkm 244.5). *Vemco Vr2* single channel monitoring receivers were placed in Deep Creek, Smith Creek and Boundary Creek in Idaho, and in the Goat River and Summit Creek in Canada to determine burbot use of the tributary streams. *Vemco* receivers in the Kootenai River were also monitored for burbot movement. Some of these transmitters carried into the winter of 2006-2007, but all batteries expired eventually. Because only two burbot were captured during winter of 2006-2007, no more fish were tagged with transmitters and only three during 2007-2008, thus it is unlikely any more will be tagged until there are adequate numbers of burbot to ensure rehabilitation goals are being met.

The SOR request attempted to "cool" the Kootenai River during winter of 2007–2008 by using a selective withdrawal system at Libby Dam. Pre-spawn water temperature was above 6°C in early November 2007 but fell below that temperature after November 10 through most of December. Water temperature was usually maintained between 1 and 4°C throughout January and February 2008 and was acceptable. River water less than 6°C is important to stimulate pre-spawn migration and water temperatures <4°C are preferred for spawning (Paragamian and Wakkinen, In Press).

The results of the 2006 and 2007 fall and winter temperature operation corroborate the availability of cooler water in the forebay of the reservoir, and the ability to provide cooler release temperatures at Libby Dam (Figure 3). Reduced river volume during this time period, in conjunction with minimized release temperatures, may allow the river to cool even further as it flows downstream to the Bonners Ferry reach. The Corps of Engineers will perform temperature modeling to determine if reduction of release volume could significantly reduce water temperature in the Bonners Ferry reach of river during winter months.

Prior to Libby Dam, winter water temperature both below the current Libby Dam site and near Bonners Ferry was substantially cooler than post-dam temperatures. The Kootenai River gradually warmed slightly as it flowed downstream, whereas current conditions allow atmospheric cooling of the river, though still not to pre-dam levels (Figure 4). The committee would like to continue to investigate the possibility of influencing ambient river temperatures during the late fall and early winter migration period, as there is water available for release during this time that is substantially cooler than previous, post-dam release temperatures, and also cooler than the minimum temperature specified by the selective withdrawal temperature curve (Figure 1).

The KVRI burbot committee is pursuing methods for reintroducing burbot into the Kootenai River, and anticipates that in future years flow requests may be made to enhance spawning conditions for these fish, should they survive and persist. The committee hopes that results of seasonal temperature experiments and modeling carried out in the interim would enable the Action Agencies to implement discharge-related SOR's in future years, with less regard to seasonal conditions, should population numbers increase to a point where temperature and flow optimization would benefit the resultant migratory spawning population.




Figure 3. 2007 Libby Dam selective withdrawal gate removal, reservoir elevation, and Kootenai River flow and temperature data. The Copeland gage is downstream of Bonners Ferry, Idaho.




Figure 4. The thermograph in the Kootenai River before and after installation of Libby Dam is reversed to the point of the now-warmer river cooling with ambient air conditions as it reaches the Bonners Ferry vicinity, rather than warming.

Literature cited:

- Breaser, S.W., F.D. Stearns, M.W. Smith, R.L. West, and J.B. Reynolds. 1988. Observations of movements and habitat preferences of burbot in an Alaskan glacial river system. *Transactions of the American Fisheries Society* 117:506-509.
- Evenson, M.J. 2000. Reproductive traits of burbot in the Tanana River, Alaska. Pages 61-70, *In: V. L. Paragamian and D. W. Willis, editors. Burbot: biology, ecology, and management. American Fisheries Society, Fisheries Management Section, Publication Number 1, Bethesda, Maryland.*
- Johnson, T. 1981. Biotope changes and life cycle of *Lota lota* in the Bothnian Sea and a coastal river. *Oesterriechische Fischereiverband* 34: 6-9.
- KVRI Burbot Committee. 2005. Kootenai River/Kootenay Lake Conservation Strategy. Prepared by the Kootenai Tribe of Idaho with assistance from S. P. Cramer and Associates. 77 pp. plus appendices.
- Morrow, J.E. 1980. The freshwater fishes of Alaska. Alaska Northwest Publishing, Anchorage, Alaska.
- McCrimmon, H.R. 1959. Observations of burbot spawning in Lake Simcoe, Ontario. *Journal of Wildlife Management* 23:447-449.
- Paragamian, V.L. 2000. The effects of variable flows on burbot spawning migrations in the Kootenai River, Idaho, USA, and British Columbia, Canada. Pages 111-123, *In: V. L. Paragamian and D. W. Willis, editors. Burbot: biology, ecology, and management. American Fisheries Society, Fisheries Management Section, Publication Number 1, Bethesda, Maryland.*
- Paragamian, V. L., J. Walters, P. Rust, R. Hardy, D. Wakkinen, and D. Kedish. 2006. Kootenai River Fisheries Recovery Investigations; April-June 2006. Idaho Dept. of Fish and Game. 9pp.
- Percy, R. 1975. Fishes of the outer Mackenzie Delta. Environment Canada, Beaufort Sea Project, Technical Report No. 8, Winnipeg.
- Robins, C.R., and E.E. Duebler. 1955. The life-history and systematic status of the burbot, *Lota lota lacustris* (Walbaum), in the Susquehanna River system. New York State Museum and Science Service Circular 39, Albany.
- Schram, S.T. 2000. Seasonal movement and mortality estimates of burbot in Wisconsin waters of western Lake Superior. Pages 90-95, *In: V.L Paragamian and D.W. Willis, editors, Burbot Biology, Ecology, and Management. Publication Number 1, Fisheries Management Section of the American Fisheries Society, Spokane, Washington, USA.*